

Novel allele *elh* of the *UBP14* gene affects plant organ size via cell expansion in *Arabidopsis thaliana*.

Rakesh David^{1*}, Pei Qin Ng^{2*}, Lisa M. Smith³ and Iain R. Searle^{4§}

¹School of Agriculture, Food and Wine, The University of Adelaide, Australia

²School of Biological Sciences, The University of Adelaide, Australia

³Department of Animal and Plant Sciences, The University of Sheffield, UK


⁴School of Biological Sciences, The University of Adelaide, Australia

[§]To whom correspondence should be addressed: iain.searle@adelaide.edu.au

*These authors contributed equally.

Abstract

Plant organ size control is an essential process of plant growth and development. The regulation of plant organ size involves a complicated network of genetic, molecular interactions, as well as the interplay of environmental factors. Here, we report a temperature-sensitive hypocotyl elongation EMS-generated mutant, hereby referred to as *elongated hypocotyl under high-temperature* (*elh*). The elongated hypocotyl phenotype was prominent when the *elh* seedlings were grown at high temperature, 28°C, but not under the growth temperature of 21°C. We observed significantly larger organ sizes in *elh* plants, including cotyledons, petals and seeds. In *elh* plants, the cell sizes in cotyledons and petals were significantly larger than wild type. By measuring the cell density and organ area of cotyledons, petals and mature dissected embryos, we found no differences in total cell numbers in any organ indicating that cell expansion rather than cell proliferation was perturbed in *elh*. *elh* plants produced leaves at a slower rate than wild type plants, suggesting that perturbing the balance between cell division and cell expansion is linked to the developmental rate at which leaves are produced.






Figure 1. The *elh* mutation affects cell and organ growth and is an allele of *UBP14*: A. Hypocotyl elongation phenotypes of wild type and *elh* *Arabidopsis thaliana* seedlings grown at 21°C and 28°C. No hypocotyl length difference between wild type and *elh* seedlings was observed when grown at 21°C, however, the *elh* hypocotyls were significantly longer when grown at 28°C (Student's t-test $p < 0.05$). B. Comparison of organ sizes (cotyledons, flowers, and seeds) of wild type and *elh*. C. Differential interference contrast (DIC) microscopy images showing cell sizes of cotyledons, petals and mature dissected embryos of wild type and *elh*. The cell density was calculated for each organ analysed. The cell densities between wild type and *elh* were significantly different in cotyledons and petals ($p < 0.05$) but not significantly different in embryos (Student's t-test $p > 0.05$). D. Leaf growth of wild type and *elh* plants. The mutant produced fewer leaves than wild type after 16 days. Minor leaf photobleaching in *elh* is caused by the presence of an unlinked transgene (Smith *et al.*, 2007) that does not affect leaf growth rate (Searle *et al.*, 2010). The unlinked transgene was removed for subsequent experiments by backcrossing to wild type Col-0 three times and then self-pollinating twice. E. The effect of naphthylphthalamic acid (NPA) on hypocotyl length in wild type and *elh* at 28°C. F. Exogenous auxin (IAA) effect on hypocotyl length in wild type and *elh* at 28°C. G. Bulk segregant analysis using a next generation mapping (NGM) approach mapped *elh* to chromosome 3. H. Transgene complementation of leaf growth rate in *elh* by using a wild type *UBP14* gene (Student's t-test, $p < 0.05$). I. Multiple sequence alignments of *UBP14* of *Arabidopsis thaliana* (At, AT3G20630), *Homo sapiens* (Hs, P54578), *Dictyostelium discoideum* (Dd, P54201), and *Saccharomyces cerevisiae* (Sc,

P38237) adapted from Doeling *et al.* (2001). The red arrow indicates the site of the non-synonymous amino acid substitution in *elh* from cysteine to tyrosine. In panels A-H, the mean and standard deviation are shown (n= 20 plants).

Description

Plant organ size control is an essential process of plant growth and development. The regulation of plant organ size involves a complicated network of genetic, molecular interactions, as well the interplay of environmental factors (Hepworth and Lenhland, 2014). Two elements of plant organ size development are cell proliferation, resulting in increase in cell numbers, and cell expansion, which increases the overall organ size via larger cell area occupancy. However, our understanding of the fine-tuned regulation of organ size remains incomprehensive. Here, we report a temperature-sensitive hypocotyl elongation EMS-generated mutant, hereby referred to as *elongated hypocotyl under high-temperature (elh)*. The elongated hypocotyl phenotype is prominent when the *elh* seedlings are grown at high temperature, 28°C, but not under the growth temperature of 21°C (Figure 1A). As hypocotyl elongation is driven primarily by cell elongation processes (Derbyshire *et al.* 2007), we compared the difference of organs size between in wild type and *elh* plants. We observed significantly larger organ sizes in *elh* plants, including cotyledons, petals and seeds when compared to the control (Student's t-test $p < 0.05$, Figure 1B). We further inspected the cell size and number of these three organs in wild type and *elh* plants. In *elh*, the cell sizes in cotyledons and petals were significantly larger than wild type (Student's t-test $p < 0.05$, Figure 1C). We also measured the cell density and organ area of cotyledons, petals and mature dissected embryos. By extrapolating cell numbers based on organ size, we found no differences in total cell numbers in either organ (Student's t-test $p > 0.05$) indicating that cell expansion rather than cell proliferation was perturbed in *elh*. Plastochron, leaf initiation rate, has been previously linked to cell expansion (Wang *et al.* 2008). We observed reduced leaf number in *elh* plants when compared to wild type (Student's t-test $p < 0.05$, Figure 1D) and a comprehensive analysis showed that *elh* plants produced leaves at a slower rate than wild type plants (Student's t-test $p < 0.05$, Figure 1H), suggesting that perturbing the balance between cell division and cell expansion is linked to the developmental rate at which leaves are produced.

As auxin is a key hormone in regulating hypocotyl elongation and cell expansion (Gray *et al.* 1998; Chapman *et al.* 2012) and high auxin levels were shown to increase hypocotyl elongation (Wang *et al.* 2008; Zhao *et al.* 2001), we hypothesised that higher auxin levels in *elh* seedlings may cause the temperature-sensitive hypocotyl elongation phenotype. To test this hypothesis, we introduced the auxin inhibitor 1-*N*-naphthylphthalamic acid (NPA) into both wild-type and *elh* seedlings (Figure 1E). Higher levels of NPA reduced the hypocotyl length in both wild type and *elh* seedlings when grown at 28°C. At the concentration of 1 μM NPA, the length of *elh* hypocotyls were similar to wild type with no NPA (Figure 1 E). When we applied exogenous auxin, indole-3-acetic acid (IAA) to wild type and *elh* seedlings (Figure 1F), the hypocotyl length elongation response at 28°C was similar in both genotypes (Figure 1F). Together these results suggest that *elh* accumulates higher auxin levels leading to a longer hypocotyl when compared to the wild type when grown at 28°C.

To understand the underlying molecular cause of the observed *elh* phenotypes, we performed bulk segregant analysis combined with Next Generation Mapping (NGM) to identify the causative mutation. Our NGM analysis suggested that the causative mutation in *elh* resulted in a non-synonymous amino acid substitution from cysteine to tyrosine in the *Ubiquitin carboxyl-terminal hydrolase 14 (UBP14; AT3G20630)* gene (Figure 1G). *UBP14* genetically complemented both the cotyledon size and plastochron phenotypes in transgenic *elh* plants (Student's t-test $p < 0.05$, Figure 1H) demonstrating that *UBP14* and *ELH* are the same gene. *UBP14* is essential for early embryo development in *Arabidopsis thaliana* as null *ubp14* mutants are embryo lethal (Doelling *et al.* 2001). Another *UBP14* allele has been shown to regulate organ size via regulating cell cycle kinases which affect both cell numbers and cell expansion (Xu *et al.* 2016). The *UBP14* allele, referred to as *da3*, has altered mRNA splicing resulting in a premature stop codon and the encoded protein lacks the conserved C-terminal His Box. Here, we characterized a novel *UBP14* allele gene that regulates cell organ size and elongation of hypocotyl under high temperature in *Arabidopsis*. Based on our findings, we postulate that *UBP14* regulates organ size via cell expansion through increased levels of auxin that affects the stability of HEAT SHOCK PROTEIN 90 (HSP90) dependent TIF1 co-receptor complexes. It has been shown that high temperatures lead to upregulation of auxin synthesis, followed by the activation of HSP90 to stabilize the TIR1 Auxin/IAA repressor degradation complex (Wang *et al.* 2016). The *elh* allele may act to destabilise the HSP90-TIR1 complex, leading to longer hypocotyls (Yamada *et al.* 2009; Wang *et al.* 2016). Unlike the *da3* mutant allele that perturbs both cell expansion and proliferation (Xu *et al.* 2016), our findings suggest that these two functions can be separated by altering protein function. The *elh* allele identified here is potentially a useful tool to understand the role of *UBP14* in plant development.

Methods

[Request a detailed protocol](#)

Plant materials and growth

Arabidopsis thaliana (Columbia accession, Col-0) wild type and mutant plants were grown in Phoenix Biosystems chambers under metal halide lights as previously described in David *et al.* (2017). For plate experiments, seeds were first

surface sterilized, plated on ½ MS medium supplemented with 1% sucrose and sealed as previously described in David *et al.* (2017). All plants were grown under long-day photoperiod conditions of 16 h light and 8 h darkness. For the temperature-sensitive hypocotyl length experiments, seedlings were grown at either 21°C or 28°C.

Auxin inhibitor NPA and exogenous auxin IAA treatments

For NPA (Sigma Cat no- 33371-100MG) treatment, wild type and *elh* seeds were germinated on ½ MS media containing various concentration of NPA and along with control plates were transferred at 1 day after germination to 28°C. Seedlings were grown for 7 days and hypocotyl length was measured as described in Gray *et al.* (1998).

For IAA (Sigma Cat no-I-5148) treatment, seeds were plated on ½ MS media without IAA. Five days post germination, seedlings were transferred to plates containing different concentrations of IAA and allowed to grow for 2 days before hypocotyl length was measured as described in Chapman *et al.* (2012).

Bulk segregant analysis and next-generation sequencing

To perform a bulk segregant analysis approach, *elh* was crossed to a polymorphic *Landsberg erecta* (*Ler*) parent and in the F₂ population, plants that had reduced plastochron were bulked together and sequenced on the Illumina HiSeq platform. The sequencing reads were trimmed using TrimGalore! (<https://github.com/FelixKrueger/TrimGalore>), trimmed reads were aligned to the TAIR10 reference genome using BWA (Li and Durbin 2010), followed by variant calling using SAMtools (Li *et al.* 2009). Identification of a SNP deserted region corresponding to *ELH* was achieved by plotting SNP frequencies across the 5 chromosomes using the Next Generation EMS mutation mapping tool (<http://bar.utoronto.ca/ngm/cgi-bin/emap.cgi>).

Genetic complementation construct and generation of transgenic plants

For the *ELH/UBP14* (AT3G20630) complementation construct, the full-length genomic sequence of *ELH/UBP14* including 2 kb upstream promoter sequence was PCR amplified and cloned into the Gateway cloning vector PCR8 TOPO-TA (Invitrogen). The respective insert was sequenced, then cloned into the destination vector pMDC100 via the Gateway Cloning approach (Curtis and Grossniklaus 2003). The construct was transformed into *elh* plants via *Agrobacterium tumefaciens*-mediated floral dipping as described by Davis *et al.* (2009). Transgenic plants were selected on kanamycin containing plates.

Acknowledgments: We thank the ACRF Cancer Genomics (Adelaide, Australia) staff for assistance with the genome sequencing.

References

- Chapman EJ, Greenham K, Castillejo C, Sartor R, Bialy A, Sun TP, Estelle M. 2012. Hypocotyl transcriptome reveals auxin regulation of growth-promoting genes through GA-dependent and -independent pathways. *PLoS One* 7: e36210. PMID: 22590525.
- Curtis MD, Grossniklaus U. 2003. A gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiol* 133: 462-9. PMID: 14555774.
- Davis AM, Hall A, Millar AJ, Darrach C, Davis SJ. 2009. Protocol: Streamlined sub-protocols for floral-dip transformation and selection of transformants in *Arabidopsis thaliana*. *Plant Methods* 5: 3. PMID: 19250520.
- David R, Burgess A, Parker B, Li J, Pulsford K, Sibbritt T, Preiss T, Searle IR. 2017. Transcriptome-Wide Mapping of RNA 5-Methylcytosine in *Arabidopsis* mRNAs and Noncoding RNAs. *Plant Cell* 29: 445-460. PMID: 28062751.
- Derbyshire P, Findlay K, McCann MC, Roberts K. 2007. Cell elongation in *Arabidopsis* hypocotyls involves dynamic changes in cell wall thickness. *J Exp Bot* 58: 2079-89. PMID: 17470442.
- Doelling JH, Yan N, Kurepa J, Walker J, Vierstra RD. 2001. The ubiquitin-specific protease UB14 is essential for early embryo development in *Arabidopsis thaliana*. *Plant J* 27: 393-405. PMID: 11576424.
- Gray WM, Ostin A, Sandberg G, Romano CP, Estelle M. 1998. High temperature promotes auxin-mediated hypocotyl elongation in *Arabidopsis*. *Proc Natl Acad Sci U S A* 95: 7197-202. PMID: 9618562.
- Hepworth J, Lenhard M. 2014. Regulation of plant lateral-organ growth by modulating cell number and size. *Curr Opin Plant Biol* 17: 36-42. PMID: 24507492.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup.. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078-9. PMID: 19505943.
- Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26: 589-95. PMID: 20080505.

Searle IR, Pontes O, Melnyk CW, Smith LM, Baulcombe DC. 2010. JM14, a JmjC domain protein, is required for RNA silencing and cell-to-cell movement of an RNA silencing signal in Arabidopsis. *Genes Dev* 24: 986-91. PMID: 20478993.

Smith LM, Pontes O, Searle I, Yelina N, Yousafzai FK, Herr AJ, Pikaard CS, Baulcombe DC. 2007. An SNF2 protein associated with nuclear RNA silencing and the spread of a silencing signal between cells in Arabidopsis. *Plant Cell* 19: 1507-21. PMID: 17526749.

Wang JW, Schwab R, Czech B, Mica E, Weigel D. 2008. Dual effects of miR156-targeted SPL genes and CYP78A5/KLUH on plastochron length and organ size in Arabidopsis thaliana. *Plant Cell* 20: 1231-43. PMID: 18492871.

Wang R, Zhang Y, Kieffer M, Yu H, Kepinski S, Estelle M. 2016. HSP90 regulates temperature-dependent seedling growth in Arabidopsis by stabilizing the auxin co-receptor F-box protein TIR1. *Nat Commun* 7: 10269. PMID: 26728313.

Xu Y, Jin W, Li N, Zhang W, Liu C, Li C, Li Y. 2016. UBIQUITIN-SPECIFIC PROTEASE14 Interacts with ULTRAVIOLET-B INSENSITIVE4 to Regulate Endoreduplication and Cell and Organ Growth in Arabidopsis. *Plant Cell* 28: 1200-14. PMID: 27099260.

Yamada M, Greenham K, Prigge MJ, Jensen PJ, Estelle M. 2009. The TRANSPORT INHIBITOR RESPONSE2 gene is required for auxin synthesis and diverse aspects of plant development. *Plant Physiol* 151: 168-79. PMID: 19625638.

Zhao Y, Christensen SK, Fankhauser C, Cashman JR, Cohen JD, Weigel D, Chory J. 2001. A role for flavin monooxygenase-like enzymes in auxin biosynthesis. *Science* 291: 306-9. PMID: 11209081.

Funding: This research was supported by an Australian Research Council Future Fellowship (FT130100525) awarded to IRS.

Author Contributions: Rakesh David: Investigation, Visualization, Writing - review and editing. Pei Qin Ng: Formal analysis, Writing - original draft, Writing - review and editing, Visualization. Lisa M. Smith: Writing - review and editing, Investigation. Iain R. Searle: Conceptualization, Investigation, Writing - original draft, Writing - review and editing, Visualization, Funding acquisition, Project administration, Supervision.

Reviewed By: Stephan Wenkel

History: Received February 12, 2021 **Revision received** May 1, 2021 **Accepted** May 28, 2021 **Published** June 1, 2021

Copyright: © 2021 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: David, R; Ng, PQ; Smith, LM; Searle, IR (2021). Novel allele *elh* of the *UBP14* gene affects plant organ size via cell expansion in *Arabidopsis thaliana*.. microPublication Biology. <https://doi.org/10.17912/micropub.biology.000401>